Referral Form Positive Fecal Immunochemical Testing

Patients must be 18 years of age at time of referral.

	ile ble Forderer wist that we of some FIT we sitis	
The target wait time from FIT postive result	ailable Endoscopist that performs FIT positive to colonscopy is 8 weeks.	colonoscopies.
-	osest to Home 🛛 Closest to Primary Care P	rovider
PATIENT INFORMATION	Gender:	
Last name:	First name:	
DOB: DD/MM/YYYY	OHIP#:	VC:
Phone:	Alternate Phone:	
Address:	City:	Postal Code:
Email:		
	ease arrange to bring a family member and,	or friend to translate for you if needed.
Barriers to Communication:	npairment 🗆 Hearing Impairment 🗆 Sight I	mpairment 🛛 Other
LAB REPORT		
Please attach FIT Positive report.		
MEDICAL HISTORY (please attach cumul	ative patient profile of medical history)	
Diabetes Sleep Apnea	Image: Morbid Obesity (BMI >40)	Asthma/COPD requiring hospitalization
CHF Bleeding Disorder Circle asian Atrial Silvaillation		□ Renal Insufficiency (eGFR)
Cirrhosis Atrial Fibrillation	PCI/Stenting in the last 12 months	Image: MI/Angina within last 12 months
MEDICATIONS	e/Dose/Frequency	ot take any medications
Is the patient on anti-coagulant medication		
If Yes, Name & Dose	Reason for treatment	
Expected Duration		
IMPORTANT INFORMATION		
Your patient will be contacted directly by the instructions.	e hospital to book his/her colonoscopy date a	nd will be provided with preparation
You will receive a consult note upon complete	tion of the procedure.	
		ith you 10 days prior to his/her colonoscopy to
review the anticoagulant management plan	. See back of referral form for guideline.	
I hereby refer the above noted patient to a	physician specialist for follow-up related to	a FIT positive report.
Primary Care Provider Name:		
REFERRING PROVIDER INFORMATION	N: 🗆 MD 🗆 NP	
Name (Printed):	Phone:	
Address:	_	
Signature:	Billing#:	Date: <u>DD/MM/YYYY</u>
ONLY CO	MPLETED REFERRALS WILL B	E ACCEPTED
	Halton	

Anticoagulant Management Guidelines				
Medication	Guideline			
ASA	Do not Hold			
Plavix, Ticagrelor	Hold for 5 days (unless recent stent or other cardiac risk)			
Coumadin	Hold for 5 days (may need bridging with LMWH if mechanical valve or high risk)			
NOAC	Hold for 2 days if normal renal function Hold for 4 days if poor EGFR with caveats as for Coumadin			
<u>For furt</u>		see: https://www.giejournal.org/article/	<u>S0016-5107(15)02950-8/pdf</u>	
Halton Healthcare (GH) Georgetown Hospital		Halton Healthcare (MDH) Milton District Hospital	Halton Healthcare (OTMH) Oakville Trafalgar Memorial Hospital	
1 Princess Anne	•	725 Bronte Street South	3001 Hospital Gate	